

FAMILY NAME

MRN

GIVEN NAME

MALE FEMALE

D.O.B. ____/____/____

M.O.

Facility:

ADDRESS



PAEDIATRIC SEPSIS PATHWAY

RECOGNISE • RESUSCITATE • REFER

LOCATION / WARD

COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE

Sepsis recognition

Date: ____/____/____ Time: ____:____

Emergency Department

Triage category 1 2 3 4 5

Inpatient Ward: _____

Clinical Review

Rapid Response



RESUSCITATE

A

Airway - Assess and maintain patent airway

B

Breathing - Assess and administer oxygen if required; aim SpO₂ ≥ 95%

C

Circulation - Vascular access, bloods, antibiotics and fluid resuscitation

Consider intraosseous access after two failed attempts at cannulation

Blood Culture(s)

Glucose Result ____ mmol/L

Lactate Result ____ mmol/L *Lactate ≥ 2mmol/L is significant*

FBC Coags LFTs EUC CRP/Procalcitonin (PCT)

Antibiotics - Prescribe and administer antibiotics within 60 minutes of sepsis recognition

Use *Therapeutic Guidelines: Antibiotic* OR locally endorsed antibiotic prescribing guideline

For neonates use *Therapeutic Guidelines: Antibiotic* and seek expert advice

First/new antibiotic administered Date: ____/____/____ Time: ____:____

Fluid Resuscitation

(intravenous or intraosseous)

Give initial 20mL/kg 0.9% sodium chloride bolus STAT

Repeat 20mL/kg 0.9% sodium chloride bolus if no improvement in heart rate, capillary refill, colour

Consider commencement of vasopressors and discuss need for intubation with senior clinician

D

Disability - Assess level of consciousness (LOC) using Alert, Voice, Pain, Unresponsive (AVPU)

E

Exposure - Targeted history and re-examine the patient for source of sepsis
Consider cerebrospinal fluid, urine, swab collection, viral culture, x-ray

F

Fluid - Monitor/document strict fluid input/output and consider IDC

G

Check Blood Glucose Level - If less than 3mmol/L treat with 2mL/kg 10% Glucose

Holes Punched as per AS2828.1: 2019
BINDING MARGIN - NO WRITING



SMR060399



SMR060399

Holes Punched as per AS2828.1: 2019

BINDING MARGIN - NO WRITING

NH700131 230920

FAMILY NAME		MRN
GIVEN NAME		<input type="checkbox"/> MALE <input type="checkbox"/> FEMALE
D.O.B. ____/____/____		M.O.
ADDRESS		
LOCATION / WARD		
COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE		

Facility:



RECOGNISE · RESUSCITATE · REFER

PAEDIATRIC SEPSIS PATHWAY

RESUSCITATE

Monitor and Reassess

Continue monitoring, assess for signs of deterioration and escalate as per local paediatric CERS

- Tachypnoea (Red or Yellow Zone)
- Persistent tachycardia (Red or Yellow Zone), slow capillary refill and hypotension
- Colour pale and mottled
- Drowsiness or decreased level of consciousness
- Urine output < 1 mL/kg/hour
- Acidosis, increasing serum lactate or procalcitonin
- Hypoglycaemia, leukopaenia or abnormal coagulation
- Consider other causes of deterioration

REFER

If no improvement Paediatric Intensive Care may be required

- Seek advice immediately from local/regional paediatric experts and/or NETS using ISBAR Tel: 1300 36 2500
- Administer further fluid bolus
- Consider second vascular access, vasopressors and intubation
- Update the Attending Medical Officer on the patient's condition using ISBAR
- Discuss the management plan with the patient and their family/carers
- Sepsis management plan documented by a medical officer in the health care record as per page 4 (over)

Name: _____ Designation: _____ Signature: _____

NO WRITING

FAMILY NAME

MRN

GIVEN NAME

 MALE FEMALE

D.O.B. ____ / ____ / ____

M.O.

Facility:

ADDRESS



RECOGNISE · RESUSCITATE · REFER

PAEDIATRIC SEPSIS PATHWAY

LOCATION / WARD

COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE

SEPSIS MANAGEMENT PLAN

Patients with presumed sepsis are at a high risk of deterioration despite initial resuscitation with intravenous antibiotics and fluids. These patients require a management plan which needs to be discussed with the Attending Medical Officer (AMO). The Infectious Diseases Physician/Clinical Microbiologist and Antimicrobial Stewardship (AMS) team are to be consulted where necessary. This plan needs to be communicated to the Senior Medical Officer, Nurse in Charge, patient and patient's family/carers.

Specific management plans are to be documented in the health care record

Initial 24 hours	Continue monitoring	<ul style="list-style-type: none"> Prescribe the frequency of observations Minimum recommendation every 30 minutes for 2 hours, then hourly for 4 hours Monitor and reassess for signs of deterioration which may include one or more of the following: <div style="background-color: #e0e0e0; padding: 5px; margin-top: 5px;"> Tachypnoea (Red or Yellow Zone) Persistent tachycardia (Red or Yellow Zone), slow capillary refill and hypotension Decreased or no improvement in level of consciousness Urine output less than 1mL/kg/hour over 4 hours No improvement in serum lactate level </div> <p>If deteriorating (has any Red or Yellow Zone criteria), escalate as per local CERS and inform AMO</p>	<input type="checkbox"/>
	Repeat lactate 4 and 8 hours post recognition	4 hours Date: ____ / ____ / ____ Time: ____ : ____ Result ____ . ____ mmol/L <input type="checkbox"/> 8 hours Date: ____ / ____ / ____ Time: ____ : ____ Result ____ . ____ mmol/L <input type="checkbox"/>	<input type="checkbox"/>
	Fluid resuscitation	<ul style="list-style-type: none"> Prescribe IV fluids as appropriate based on the patient's condition <i>Monitor for signs of fluid overload/pulmonary oedema/inappropriate antidiuretic hormone</i> 	<input type="checkbox"/>
	Reassess	<ul style="list-style-type: none"> Confirm diagnosis and consider other causes of deterioration e.g. dehydration/hypovolaemia/haemorrhage or an overdose/over sedation Check preliminary results <i>If patient is neutropenic, review antibiotics and change if required</i> 	<input type="checkbox"/>
	Review treatment/management	<ul style="list-style-type: none"> Discuss with AMO Document plan to continue, change or cease antibiotics Continue monitoring for deterioration including urine output If the patient's recovery is uncertain discuss the goals of care with the patient's family/carers 	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
24 - 48 hours	Reassess	<ul style="list-style-type: none"> Actively seek microbiology/investigation results and review Confirm diagnosis, document source of sepsis in the health care record Discuss with AMO Consider seeking advice from infectious disease/microbiology physician Document plan to continue, change or cease antibiotics Obtain AMS approval for restricted antibiotics Repeat biochemistry as indicated Continue monitoring for deterioration including urine output 	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
Continue to monitor as per patient's condition – observations, medical review, antibiotics			

Holes Punched as per AS2828.1: 2019
BINDING MARGIN - NO WRITING

SMR060399

