

NEWBORN ANTIBIOTIC GUIDELINE

for early and late onset sepsis
during birth episode of care

Revised June 2018



© Clinical Excellence Commission 2018

All rights are reserved. In keeping with the NSW Government's commitment to encouraging the availability, dissemination and exchange of information (and subject to the operation of the *Copyright Act 1968*), you are welcome to reproduce the information which appears in this publication, as long as the user of the information agrees to:

- use the document for information only
- save or print a single copy for personal use only and not to reproduce any major extract or the entire document except as permitted under *Copyright Act 1968* (as amended) without the prior written permission of the State of New South Wales
- acknowledge the source of any selected passage, table diagram or other extract reproduced
- not make any charge for providing the Information to another person or organisation without the prior written consent of the State of New South Wales and payment of an agreed copyright fee
- not modify the Information without the express prior written permission of the State of New South Wales
- include this copyright notice in any copy made:

© - Copyright – Clinical Excellence Commission for and on behalf of the Crown in right of the State of New South Wales.

Source **antibacterial recommendations, Neonatal Medicines Formulary Consensus Group with permission.**

<https://www.seslhd.health.nsw.gov.au/royal-hospital-for-women/australasian-neonatal-medicines-formulary-anmf>

Accessed on 10 April 2018

National Library of Australia Cataloguing-in-Publication entry

Title: Newborn Antibiotic Guideline for early and late onset sepsis during birth episode of care

SHPN: (CEC) 180357

Suggested citation

Clinical Excellence Commission, 2018, Newborn Antibiotic Guideline for early and late onset sepsis during birth episode of care. Revised June 2018. Sydney: Clinical Excellence Commission

Clinical Excellence Commission

Board Chair: Associate Professor Brian McCaughan, AM

Chief Executive: Ms. Carrie Marr

Any enquiries about or comments on this publication should be directed to:

Clinical Excellence Commission

Locked Bag 8

Haymarket NSW 1240

Phone: (02) 9269 5500

Email: cec-sepsis@health.nsw.gov.au

CONTENTS

Document purpose	4
Important points to consider when using this guideline	6
Decision tree	7
Early onset of sepsis associated with birth (< 72 hours old)	8
Late onset of sepsis associated with birth (\geq 72 hours old)	9
Medication administration table	10
References and bibliography	13

DOCUMENT PURPOSE

The Clinical Excellence Commission's (CEC) *Newborn Antibiotic Guideline for early and late onset sepsis during birth episode of care* aims to guide the prescription and timely administration of antibiotics for **newborns during the birth episode of care**.

Definitions of sepsis^{1, 2, 3}

SEPSIS - Early onset

Early-onset sepsis < 72 hours of age is associated with acquisition of microorganisms from the mother. Most common microorganisms associated with early-onset infection include the following:

- Group B Streptococcus (GBS)
- *Escherichia coli*
- *Haemophilus influenzae*
- *Listeria monocytogenes*
- Herpes simplex virus

SEPSIS - Late onset

Late-onset sepsis occurs \geq 72 hours of age and may also be acquired from the caregiving environment. Organisms that have been implicated in causing late-onset sepsis include the following:

- Group B Streptococcus
- *Escherichia coli*
- *Staphylococcus aureus*
- Other gram negative organisms
- Herpes simplex virus
- Coagulase-negative staphylococci


This guideline is not intended for:

- Neonates who present to an Emergency Department after birth → use *Paediatric Antibiotic Guideline for Severe Sepsis and Septic Shock and Unwell Neonates*
- Premature neonates being managed in specialised Neonatal Intensive Care Units (NICUs) with locally-endorsed antibiotic guidelines
- Paediatric patients (29 days – 16 years of age) → use **Paediatric Antibiotic Guideline for Severe Sepsis & Septic Shock and Unwell Neonates**

IMPORTANT POINTS TO CONSIDER WHEN USING THIS GUIDELINE

- The selection of appropriate antibiotic therapy is complex - this guideline is not intended to cover all possible scenarios
- Prompt administration of antibiotics and resuscitation fluids is vital in the management of the newborn with sepsis. In newborns diagnosed with early or late onset of sepsis, the goal is to commence antibiotic therapy within the first hour
- Obtain at least one set of blood cultures and other clinical specimens (e.g. urine, cerebrospinal fluid, wound swabs) as appropriate **PRIOR TO** antibiotic commencement where possible - do not delay antibiotic administration if a blood culture cannot be obtained or to wait for results of investigations
- If agents listed here are not available in your hospital, consult the Attending Medical Officer and seek expert advice
- Newborn patients must be bare weighed to ensure correct calculation of medication dosing – birth weight is to be used for calculation of medication dosing until the newborn crosses birth weight
- Clinicians must document the indication, drug name, dose, route of administration and review date for antibiotics in the newborn's health care record
- The neonate and antibiotic treatment within 24 and 48 hours or once microbiology results are available and antibiotics continued, changed or ceased as required. Where necessary, discuss with the Paediatrician, Neonatologist or Clinical Microbiologist

NEWBORN ANTIBIOTIC GUIDELINE FOR EARLY AND LATE ONSET SEPSIS DURING BIRTH EPISODE OF CARE DECISION TREE



Further management:

The newborn with presumed or suspected sepsis must be discussed with a consultant Paediatrician or Neonatologist. If a specialist is not available, call NETS NSW phone **1300 36 2500** for urgent advice.

The newborn must be reviewed by the Attending Medical Officer within 24 - 48 hours of commencing the sepsis pathway and antibiotic therapy, with referral to the infectious diseases and/or clinical microbiology service for specific advice if required. The management plan should be communicated to the Senior Medical Officer, Midwife/Nurse in Charge, and the newborn's family/carers.

Clinicians who are experiencing difficulty in interpreting microbiology results when rationalising antibiotic therapy should contact the designated infectious diseases and/or clinical microbiology service.

INDICATION: EARLY ONSET SEPSIS ASSOCIATED WITH BIRTH (< 72 hours old)

ROUTE OF ADMINISTRATION

INTRAVENOUS (IV)	INTRAMUSCULAR (IM) <i>Should only be used in the short term until IV access established</i>
benzylpenicillin 60 mg/kg, 12-hourly	benzylpenicillin 60 mg/kg, 12-hourly
PLUS	PLUS
gentamicin 5 mg/kg, daily [Note 1]	gentamicin 5 mg/kg, daily [Note 1]
OR	OR
ampicillin 50 mg/kg, 12-hourly	ampicillin 50 mg/kg, 12-hourly
PLUS	PLUS
gentamicin 5 mg/kg, daily [Note 1]	gentamicin 5 mg/kg, daily [Note 1]
<i>If newborn is severely unwell with suspected or known meningitis ADD</i> cefotaxime 50 mg/kg, 8-hourly	<i>If newborn is severely unwell with suspected or known meningitis ADD</i> cefotaxime 50 mg/kg, 8-hourly
<i>If Herpes simplex infection is known or suspected ADD</i> aciclovir 20 mg/kg, 8-hourly	<i>If Herpes simplex infection is known or suspected</i> SEEK EXPERT ADVICE

Note 1: Refer to Neonatal Medicines Formulary Consensus Group <https://www.seslhd.health.nsw.gov.au/royal-hospital-for-women/australasian-neonatal-medicines-formulary-anmf> for ongoing monitoring

INDICATION: LATE ONSET SEPSIS ASSOCIATED WITH BIRTH (≥ 72 hours old)

ROUTE OF ADMINISTRATION

INTRAVENOUS (IV)	INTRAMUSCULAR (IM) <i>Should only be used in the short term until IV access established</i>
flucloxacillin 50 mg/kg 0-7 days old, 12-hourly 8-28 days old, 8-hourly PLUS	flucloxacillin 50 mg/kg 0-7 days old, 12-hourly 8-28 days old, 8-hourly PLUS
gentamicin 5mg/kg, daily If newborn has suspected or known meningitis REPLACE flucloxacillin with: ampicillin 50 mg/kg 0-7 days old, 12-hourly 8-28 days old, 6-hourly PLUS	gentamicin 5mg/kg, daily If newborn has suspected or known meningitis REPLACE flucloxacillin with: ampicillin 50 mg/kg 0-7 days old, 12-hourly 8-28 days old, 6-hourly PLUS
cefotaxime 50 mg/kg 0-7 days old, 8-hourly 8-28 days old, 6-hourly PLUS	cefotaxime 50 mg/kg 0-7 days old, 8-hourly 8-28 days old, 6-hourly PLUS
gentamicin 5 mg/kg, daily [Note 1] If Herpes simplex infection is known or suspected ADD aciclovir 20 mg/kg, 8-hourly If MRSA is known or suspected ADD vancomycin 15 mg/kg (maximum single dose, 90 mg) 0-7 days old, 12-hourly 8-28 days old, 8-hourly If abdominal sepsis is known or suspected ADD metronidazole 15 mg/kg as a loading dose then subsequent doses of: 7.5 mg/kg 0-7 days old, 8-hourly 7.5 mg/kg 8-28 days old, 6-hourly	gentamicin 5 mg/kg, daily [Note 1] If Herpes simplex infection is known or suspected SEEK EXPERT ADVICE If MRSA is known or suspected SEEK EXPERT ADVICE SEEK EXPERT ADVICE

Note 1: Refer to Neonatal Medicines Formulary Consensus Group <https://www.seslhd.health.nsw.gov.au/royal-hospital-for-women/australasian-neonatal-medicines-formulary-anmf> for ongoing monitoring

MEDICATION ADMINISTRATION TABLE Adapted with permission from Neonatal Medicines Formulary Consensus Group

- From a microbiological perspective, injectable medication **must be prepared immediately prior to administration** using aseptic technique
- Reconstitute antibiotics with sterile water for injection unless stated otherwise in the table below
- Displacement volume is the volume that the powder component of a drug takes up upon reconstitution. It needs to be added to the diluent volume to ensure accuracy when calculating doses that are less than a full vial. Thus the diluent volume recommended in the Product Information (PI) may sometimes differ from the volume recommended in this guideline. The displacement volume provided is an estimate and this may vary between brands. Please check in the Product Information or with the manufacturer

volume of diluent to reconstitute a vial + displacement volume of drug powder = final volume of vial

- If further dilution is required for IV injection or infusion, use sterile sodium chloride 0.9% or sterile glucose 5% unless stated otherwise.
- Where possible use separate dedicated lines for resuscitation fluid and for medications. When injecting antibiotics directly into an IV injection port which has resuscitation fluid running:
 - clamp the infusion fluid line and flush with 0.5 – 1mL sterile sodium chloride 0.9% solution
 - administer antibiotic over the required time
 - flush the line with 0.5 – 1 mL sterile sodium chloride 0.9% solution and recommence resuscitation fluid

Medication	Availability	Reconstitution fluid/volume	Administration	Notes
aciclovir	Vial 250 mg/10 mL	250 mg/10 mL vial: add 40 ml of water for injection to make 5 mg/mL solution	IV infusion: over 60 minutes	DO NOT GIVE INTRAMUSCULARLY Dose interval adjusted if renal impairment
ampicillin	Vial 500 mg, 1 g	500 mg vial: add 4.7 mL of water for injection to make 100 mg/mL solution 1 g vial: add 9.3 mL of water for injection to make 100 mg/mL solution IM: add 1.7 mL of water for injection to the 500 mg vial for reconstitution to make 250 mg/mL solution	IV infusion: over 5–10 minutes into the proximal cannula site with a maximum rate of 100 mg/minute	Contraindicated in patients with severe hypersensitivity to penicillins, carbapenems and cephalosporin antibiotics Separate from aminoglycosides by clearing the lines with a flush as ampicillin inactivates them Higher doses should be diluted to 30 mg/mL and infused over 30 minutes In renal impairment the excretion of ampicillin will be delayed. In infants with severe renal impairment it may be necessary to reduce the total daily dose

Medication	Availability	Reconstitution fluid/volume	Administration	Notes
benzylpenicillin	Vial 600 mg	600 mg vial: add 3.6 mL of water for injection to make 150 mg/mL solution	IV infusion: over 30 minutes	Contraindicated in patients with severe hypersensitivity to penicillins, carbapenems and cephalosporin antibiotics Separate from aminoglycoside administration by clearing the line with a flush as penicillins inactivate aminoglycosides
		IM: add 1.6 mL water for injection to the 600 mg vial to make 300 mg/mL solution	IM injection: inject deep into a large muscle	
cefotaxime	Vial 500 mg, 1 g	500 mg vial: add 4.8 mL of water for injection to make 100 mg/mL solution	IV injection: Over 3–5 minutes IV infusion: Infuse over 15–30 minutes via syringe driver	Contraindicated in patients with severe hypersensitivity to penicillins, carbapenems and cephalosporin antibiotics
		1 g vial: add 9.6 mL of water to make 100 mg/mL solution		
		IM: add 2 mL of water for injection to the 500 mg powder for reconstitution to make 230 mg/mL solution Vial can be reconstituted with lignocaine 0.5% to reduce pain of injection	IM injection: Inject deep into the gluteal muscle	
flucloxacillin	Vial 500 mg, 1 g	500 mg: add 4.6 mL of water for injection to make 100 mg/mL solution 1 g: add 9.3 mL of water for injection to make 100 mg/mL solution Draw up 2.5 mL (250 mg) and add 2.5 mL sodium chloride 0.9% to make a final volume of 5 mL with a concentration of 50 mg/mL	IV injection: Slow injection over 3–5 minutes	

Medication	Availability	Reconstitution fluid/volume	Administration	Notes
		IM: 500 mg: add 1.6 mL water for injection or lignocaine 0.5% or 1% to make a final concentration 250mg/1mL 1 g: add 3.3 mL water for injection or lignocaine 0.5% or 1% to 1 g to make a final concentration 250 mg/1mL	IM injection: Inject deep into the gluteal muscle	flucloxacillin causes significant pain and irritation with IM use
gentamicin	10 mg/1 mL ampoule – paediatric strength	10 mg/1 mL – paediatric strength: Add 1 mL (10 mg) of gentamicin to 4 mL sodium chloride 0.9% to make a final volume of 5 mL with a concentration of 2 mg/mL	IV injection: Slow infusion over 5 minutes	Gentamicin is inactivated by penicillins and cephalosporins so should not be mixed in the same solution or administered simultaneously Ensure the line is adequately flushed if administered consecutively
	80 mg/2 mL ampoule – adult strength	80 mg/2 mL – adult strength: Add 1 mL (40 mg) of gentamicin to 19 mL sodium chloride 0.9% to make a final volume of 20 mL with a concentration of 2 mg/mL		
		IM: Reconstitution not required	IM injection: administer undiluted	
metronidazole	Infusion bag 500 mg/100mL	Reconstitution not required	IV infusion: 20 - 30 minutes. Maximum rate is 25 mg/minute	DO NOT GIVE INTRAMUSCULARLY
vancomycin	Vial 500 mg, 1 g	500 mg: add 10 mL of water for injection to make a 50 mg/mL solution Draw up 1 mL (50 mg) of vancomycin and add 9 mL glucose 5% or sodium chloride 0.9% to make a final volume of 10 mL with a final concentration of 5 mg/mL	IV infusion: 60 minutes	DO NOT GIVE INTRAMUSCULARLY Monitor renal function, full blood count, hearing function and serum vancomycin concentrations Trough level: 10 - 20 mg/L (aim for higher trough level: 15 - 20 mg/L in suspected severe sepsis e.g., MRSA, bone infection, meningitis, endocarditis)

Medication	Availability	Reconstitution fluid/volume	Administration	Notes
vancomycin continued		To prepare 10 mg/mL concentration: Add 10 mL of water for injection to the 500 mg vial to make a 50 mg/mL solution Draw up 2 mL (100 mg) of vancomycin and add 8 mL glucose 5% or sodium chloride 0.9% to make a final volume of 10 mL with a final concentration of 10 mg/mL		Trough concentration should be taken within one hour prior to the 2nd dose for 18 hourly dosing and 4th dose for all other frequencies Check concentration prior to the 4th dose or after any change in dose or frequency. Perform weekly monitoring for prolonged courses More frequent monitoring may be required in renal impairment, those receiving other nephrotoxic drugs or in suspected severe sepsis

References:

1. Klinger G, Levy I, Sirota L, et al. Epidemiology and risk factors for early onset sepsis among very-low-birthweight infants. *Am J Obstet Gynecol.* 2009 Jul. 201(1):38.e1-6. [\[Medline\]](#).
2. van den Hoogen A, Gerards LJ, Verboon-Maciolek MA, Fleer A, Krediet TG. Long-term trends in the epidemiology of neonatal sepsis and antibiotic susceptibility of causative agents. *Neonatology.* 2010. 97(1):22-8. [\[Medline\]](#)
3. Lin FY, Weisman LE, Azimi P, et al. Assessment of Intrapartum Antibiotic Prophylaxis for the Prevention of Early-onset Group B Streptococcal Disease. *Pediatr Infect Dis J.* 2011 Sep. 30(9):759-763. [\[Medline\]](#).[\[Full Text\]](#)
4. Neonatal Medicines Formulary Consensus Group. <https://www.seslhd.health.nsw.gov.au/royal-hospital-for-women/australasian-neonatal-medicines-formulary-anmf> Accessed on 7 November 2017.

Bibliography

1. AMH Children's Dosing Companion accessed via CIAP <https://childrens.amh.net.au.acs.hcn.com.au/>
2. Safe Gentamicin prescribing in paediatrics. Taking the confusion and harm out of gentamicin dosing and monitoring http://www.cec.health.nsw.gov.au/_data/assets/pdf_file/0011/386291/Safe-Gentamicin-Prescribing-In-Paediatrics-Oct-2017.pdf



CLINICAL
EXCELLENCE
COMMISSION

Clinical Excellence Commission
Locked Bag 8
Haymarket NSW 1240
Phone: (02) 9269 5500
Email: CEC-SPC@health.nsw.gov.au
Web: www.cec.health.nsw.gov.au