

INFECTION CONTROL FOR FOOD SERVICE

KITCHEN

WARD

CONTACT PRECAUTIONS

DROPLET PRECAUTIONS

AIRBORNE PRECAUTIONS



Perform hand hygiene on entry



Perform hand hygiene on ward entry, exit and after touching patient surrounds



Perform hand hygiene on ward entry, exit and after touching patient surrounds



Do not enter the patient zone or approach the bedside. Perform hand hygiene on ward entry, exit.



Do not enter the patient zone or approach the bedside. Perform hand hygiene on ward entry, exit.



Standard cleaning and sanitising will destroy any germs



Order meals (paper-based and tablet) at bedside



Order meals (paper-based and tablet) at bedside



Do not enter the patient zone or approach the bedside. Take orders via phone/ non-contact method



Do not enter the patient zone or approach the bedside. Take orders via phone/ non-contact method



The dishwasher will destroy any germs



Deliver meals and collect meals/mid-meals at bedside



Deliver meals and collect meals/mid-meals at bedside



Do not enter the patient zone or approach the bedside. Deliver meals/mid-meals and collect tray/trolley from designated area



Do not enter the patient zone or approach the bedside. Deliver meals/mid-meals and collect tray/trolley from designated area



Wear gloves and an apron



Wear gloves for tray collection



Wear gloves for tray collection. For each contact precaution bed: doff gloves, perform hand hygiene and don new gloves



Wear gloves to collect trolley and pick up food tray from designated area. After collection doff gloves and perform hand hygiene



Wear gloves to collect trolley and pick up food tray from designated area. After collection doff gloves and perform hand hygiene.



Single-use wares and foam trays are not required



Single-use wares and foam trays are not required



Single-use wares and foam trays are not required



Single-use wares and foam trays are not required

Change any PPE immediately if soiled or wet/damp.

Any worker on a documented Risk Management Plan for COVID-19 or non-infectious symptom management is required to wear a mask at all times.

Where patient has a combined contact AND droplet precautions, follow droplet precautions

Where patient has a combined contact, droplet AND airborne precautions, follow airborne precautions