

CEC Post Fall Assessment and Management Guide For All Adult Patients

A fall is an event which results in a person coming to rest inadvertently on the ground or floor or other lower level: (ACSQHC 2009 Falls Best Practice Guidelines).

Please note: if a patient is found on the floor or lower level, it should be assumed that a fall has occurred unless there is reasonable evidence of a sudden onset of paralysis, epileptic seizure, loss of consciousness or overwhelming external force (being applied).

A fall event can be serious and cause injury and even death. In hospital a patient fall can be a flag that the patient's underlying medical condition may have deteriorated. The causes of falls are complex, and immediate post fall care will help to reduce the degree of harm to the patient.

As patients may have complex health issues, decisions in regards to treatment options following the fall should be discussed with the patient &/or person responsible, family/carer. In the event that a patient has an Advance Care Plan or Directive in place, appropriate symptom management will remain a priority in the plan of care.

Post fall assessment and management for all adult falls.

The post fall guide aligns with the CEC Between the Flags program. Staff are to follow local Clinical Emergency Response Systems (CERS) and if at any time a staff member and or a family member is concerned about the patient, staff are to call for a Clinical Review. Refer to: *Recognition and Management of Patients Who Are Clinically Deteriorating*, PD2011_077, 06-Dec-2011. It is assumed that staff will have undertaken DETECT (Detecting Deterioration, Evaluation, Treatment, Escalation and Communicating in Teams) training.

Immediate response: When a patient has a fall, staff are required to undertake rapid assessment and basic life support (if required) and complete observations.

Observations & ongoing monitoring: Undertake further observations and monitoring as the patient may develop symptoms later that can indicate serious changes in their condition.

Older people who fall in hospital are at risk of intracranial injury due to physiological changes in the brain. This means that patients who fall in hospital (falls and hits head, falls and does not hit head, and unwitnessed falls) are all at risk. Early signs of deterioration are fluctuating behaviours (increased agitation, restlessness or changes in alertness – lethargy, flattened) or increasing confusion. Ongoing monitoring is important.

Assessment of injury: Do not move the patient until assessed. Examine the cervical spine and immobilise if there is any indication or potential for a spinal injury. If there is any sign of injury, staff are to contact the relevant senior nurse and medical officer for a clinical review. Signs of injury may include: bruising, lacerations, swelling, redness, abrasions, shortening of limbs, restricted limb movement, external rotation of lower limbs, inability to weight bear, pain on applying pressure, and signs of deformity.

If there is a suspected head injury the decision to CT scan will depend upon the patients overall medical condition and the location and availability of appropriate services. If CT scan is not available and the patient has no other identified risk factors, clinical judgment can be used to justify continued observations rather than transfer for CT scan.

Risk factors indicating potentially significant mild head injury

- | | | |
|--|--|--|
| ■ GCS <15 at 2 hours post injury | ■ Age >65 years | ■ Large scalp haematoma or laceration ** |
| ■ Deterioration in GCS | ■ Post traumatic seizure | ■ Multi-system trauma** |
| ■ Focal neurological deficit | ■ Prolonged loss of consciousness (>5 min) | ■ Known neurosurgery / neurological deficit ** |
| ■ Clinical suspicion of skull fracture | ■ Persistent post traumatic amnesia (AWPTAS <18/18)* | ■ Delayed presentation or representation** |
| ■ Vomiting (especially if recurrent) | ■ Persistent abnormal alertness / behaviour / cognition* | * <i>particularly if persists at 4 hours post time of injury</i> |
| ■ Known coagulopathy / bleeding disorder | ■ Persistent severe headache* | ** <i>clinical judgement required</i> |

Adapted from: Algorithm 1: Initial Management of Adult Closed Head Injury: PD2012_013: Initial Management of Closed Head Injury in Adults.

Treatment must be discussed with the patient/family/carers and documented in the patient health record. Staff are to follow local health service protocols and further information can be found in the NSW Ministry of Health PD2012_013: Initial Management of Closed Head Injury in Adults.

http://www0.health.nsw.gov.au/policies/pd/2012/PD2012_013.html

The full guideline and associated algorithms provide a comprehensive guide to the management of head injury. These resources are available at: http://www.itim.nsw.gov.au/wiki/Head_injury_CPG

Assessment of risk of bleeding: A patient with known **coagulopathy** and those patients on **anticoagulant therapy** are at increased risk of bleeding and Traumatic Brain Injury (TBI) if they fall. Investigations should include a check of INR/ATTP. Continued monitoring and observation is important. If the patient presents with abnormal alertness/behaviour/cognition this may indicate subtle brain injury better than GCS and staff are to be aware and report any changes. Family/carers may help establish what is normal.

Patients at risk include:

- Patient receiving anti-coagulant therapy (e.g. warfarin, heparin, enoxaparin (Clexane), dalteparin (Fragmin), rivaroxaban),
- Patient receiving anti-platelet therapy (e.g. aspirin, clopidogrel, aspirin/dipyridamole (Assantin))
- Patient has haematological disease
- Patient has renal failure – end stage and haemodialysis patients
- Patient has chronic liver disease
- Patient is alcohol dependent

Assessment for Sepsis: A patient who falls may have sepsis. Sepsis arises when the body's response to infection causes a generalised inflammatory response. The sepsis risk factors and signs and symptoms that can contribute to the fall include: hypotension, tachypnoea, confusion, debilitated condition (frailty and poor mobility, poor balance and muscle strength) and urinary incontinence.

It is a medical emergency just like a heart attack or stroke, and can lead to shock, multiple organ failure and death, especially if not recognised early and treated promptly. Sepsis causes more deaths per year than prostate cancer, breast cancer and HIV/AIDS combined.

All patients in hospital are at risk of sepsis. Any patient with known or suspected infection or following a fall should be assessed using the Clinical Excellence Commission sepsis pathway (adult patients and paediatric).

The sepsis pathway is a tool designed to provide clear guidelines for sepsis recognition, notification, escalation and initial management.

For further information go to:

<http://www.cec.health.nsw.gov.au/programs/sepsis>

Assessment for delirium: Delirium is a clinical syndrome that may involve a disturbance in cognition, leading to changes in consciousness, attention, perception and inability to focus. Delirium is acute, developing over hours or days and can be fluctuating. A delirium can develop quickly and may be associated with extremes of behaviour (increased agitation, restlessness, changes in alertness), or delusions, hallucinations and aggression.

The Confusion Assessment Method (CAM) is the tool designed to identify and recognise delirium quickly. Staff are to complete a CAM for patients with changes in cognition and to investigate the underlying cause for delirium eg infection, possible head injury.

Staff are to continue to monitor the patient. If the patient develops fluctuating changes in cognition (increased agitation, restlessness, changes in alertness, lethargy/flattened) staff are to call for a clinical review or rapid response. A CT scan is recommended.

Assessment of ward environment and equipment: Modify any environmental risk factors that may have contributed to the fall. Equipment that has contributed to the patient's falls incident is to be reviewed and repaired or replaced if required.

Assessment of staffing levels and the need for increased patient supervision: Review the patients care plan and discuss with the Nurse Manager/Supervisor any issues related to staffing. If there is a requirement for increased patient supervision, health facilities will have in place protocols for staff to follow.

Communication and documentation: Following a fall, the patients care plan will need to be reviewed. Complete the Falls Risk Screen, Ontario Modified Stratify (SS) to ascertain falls risks and revise the patients care plan using the CEC Falls Risk Assessment and Management Plan (FRAMP) which provides a guide to suitable care interventions. Discuss the care plan with the patient and family/carer. Communicate any changes to care plan at ward handover. The Nursing Unit Manager/ Senior Staff and Medical Officer are to be notified of the fall.

A post- fall sticker may be completed and included in the patient clinical record. For serious falls incidents (SAC1 & 2) a review of the fall event with clinical leadership team will be undertaken.

All patient fall events, treatment plans and revised interventions are to be documented in the patient health record. Complete the Incident Information Management System (IIMS) notification.

For further falls information and resources which include flyers with relevant messages for patients families and carers go to:

<http://www.cec.health.nsw.gov.au/programs/falls-prevention>